

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

Doc. 1-1 on ss 2 from WPIL using MAX

©Derwent Information

## Luteinising hormone releasing hormone (LHRH) antagonists - have high anti:ovulation and low histamine releasing activity, for treatment of endometriosis, precocious puberty etc.

**Patent Number : WO9208733**

International patents classification : A61K-038/00 C07K-000/00 C07K-007/00 C07K-007/06 C07K-007/20 C07K-007/23 A01N-037/18 A61K-037/00 A61K-037/02 A61K-037/43 A61K-038/24 C07K-003/00 C07K-003/08 C07K-005/00 C07K-007/02 C07K-007/08 C07K-099/56

### • Abstract :

WO9208733 A Design and synthesis of LHRH (luteinising hormone releasing hormone) antagonists (I) comprises modifying both basic and lipophilic regions of the highly active antagonist (II) NAc-(D)-2-Nal-(D)-p-CIPhe-(D)-3-Pal-Ser-Tyr-Arg-Leu-Arg-Pro-D- Ala-NH<sub>2</sub> (II); 2-Nal=beta-(2-naphthyl)alanine; p-CIPhe = p-chloro- phenylalanine; 3-Pal = beta-(3-pyridyl)alanine so as to produce cpds. with both high anti-ovulatory activity (AOA) and low histamine-releasing activity (HRA), based on topological similarity with the neuropeptide substance P. Antagonists (I) of formula NAc-(D)-2Nal-AA-AA-Ser-AA-AA-Leu-AA-Pro-(D)-Ala-NH<sub>2</sub> (Ia) are new. In (Ia), each AA = natural or unnatural amino acid D- or L-ArAla; Ar = 1- or 2-naphthyl; H; phenyl (opt. 4-substd. by F or Cl); 2-, 3- or 4-pyrimidyl; 2-, 3- or 4-quinolyl; 1-imidazolyl; 4-pyridazyl; 5-pyrimidyl; 2-pyrazinyl; 2-oxo-1,2-dihydroquinolin-4-yl or 4-XCH<sub>2</sub>-phenyl; X = piperidino, morpholino, pyrrolidino or R<sub>2</sub>N; each R = H or 1-4C alkyl.

USE/ADVANTAGE - (I) are used to treat disorders of the reproductive endocrine system (e.g. endometriosis, precocious puberty, prostate or breast cancer); as male or female contraceptive and for diagnosis or treatment of infertility. Unlike known antagonists, they have negligible side effects.

(Dwg.0/0)

EP-564466 B (Nac-D2Nal1, DpCIPh2,Dphe3, Ser4, Arg5, DePal6, Leu7, Arg8, Pro9, DAla10)NH<sub>2</sub>. (Dwg.0/0)

### • Publication data :

Patent Family : WO9208733 A1 19920529 DW1992-24 C07K-007/20 Eng 50p \* AP: 1991WO-EP02110 19911108 DSNW: AU CA CS FI HU JP KR NO PL SU DSRW: AT BE CH DE DK ES FR GB GR IT LU NL SE

AU9188612 A 19920611 DW1992-37 C07K-007/20 FD: Based on WO9208733 AP: 1991AU-0088612 19911108; 1991WO-EP02110 19911108

ZA9108847 A 19920826 DW1992-40 C07K-000/00 48p AP: 1991ZA-0008847 19911107

PT--99458 A 19921030 DW1992-47 C07K-007/00 AP: 1991PT-0099458 19911108

CN1061605 A 19920603 DW1993-07 C07K-007/00 AP: 1990CN-0108955 19901110

NZ-240505 A 19930428 DW1993-20 C07K-007/20 AP: 1991NZ-0240505 19911107

FI9303156 A 19930709 DW1993-39 C07K-000/00 AP: 1991WO-EP02110 19911108; 1993FI-0003156 19930709

NO9301697 A 19930707 DW1993-39 C07K-007/20 AP: 1991WO-EP02110 19911108; 1993NO-0001697 19930510

**EP-564466** A1 19931013 DW1993-41 C07K-007/20 Eng

FD: Based on WO9208733 AP: 1991EP-0919435 19911108; 1991WO-EP02110 19911108 DSR: AT BE CH DE DK ES FR GB GR IT LI LU NL SE

CZ9300848 A3 19940216 DW1994-14 C07K-007/20 AP: 1993CZ-0000848 19911108

SK9300453 A3 19931006 DW1994-20 C07K-007/20 AP: 1993SK-0000453 19930510

AU-662496 B 19950907 DW1995-44 C07K-007/20 FD: Previous Publ. AU9188612; Based on WO9208733 AP: 1991AU-0088612 19911108

HU7070166 T 19950928 DW1995-46 C07K-007/20 FD: Based on WO9102110 AP: 1991WO-EP02110 19911108; 1993HU-0001353 19911108

EP-564466 B1 19970305 DW1997-14 C07K-007/23 Eng 28p FD: Based on WO9208733 AP: 1991EP-0919435 19911108; 1991WO-EP02110 19911108 DSR: AT BE CH DE DK ES FR GB GR IT LI LU NL SE

DE69125024 E 19970410 DW1997-20 C07K-007/23 FD: Based on EP-564466; Based on WO9208733 AP: 1991DE-6025024 19911108; 1991EP-0919435 19911108; 1991WO-EP02110 19911108

ES2100965 T3 19970701 DW1997-36 C07K-007/23 FD: Based on EP-564466 AP: 1991EP-0919435 19911108

**US5783562** A 19980721 DW1998-36 A61K-038/00

AP: 1991US-0789730 19911112; 1994US-0265631 19940624; 1995US-0450951 19950523

CZ-284168 B6 19980916 DW1998-43 C07K-007/23 FD:

Previous Publ. CZ9300848; Based on WO9208733 AP: 1991WO-

### • Patentee & Inventor(s) :

Patent assignee : (ASTA ) ASTA MEDICA AG  
(BIRT-) BIRTH CONTROL INST

Inventor(s) : XIAO S; SHAOBO X; XIAO ST

EP02110 19911108; 1993CZ-0000848 19911108  
RU2123499 C1 19981220 DW2000-17 C07K-007/06 AP:  
1993RU-0004994 19911108  
Priority n° : 1990CN-0108955 19901110  
Covered countries : 32  
Publications count : 19  
Cited patents : EP-277829 6.Jnl.Ref

• **Accession codes** :


Accession N° : 1992-200134 [24]  
Sec. Acc. n° CPI : C1992-091098

• **Derwent codes** :

Manual code : CPI: B04-C01C B12-E09  
B12-G01A B12-G07 B12-K03 B12-K04A  
Derwent Classes : B05

• **Update codes** :

Basic update code : 1992-24  
Equiv. update code : 1992-37; 1992-40;  
1992-47; 1993-07; 1993-20; 1993-39; 1993-  
41; 1994-14; 1994-20; 1995-44; 1995-46;  
1997-36; 1998-36; 1998-43; 2000-17



(19)

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 0 564 466 B1**

(12)

**EUROPEAN PATENT SPECIFICATION**

(45) Date of publication and mention  
of the grant of the patent:  
**05.03.1997 Bulletin 1997/10**

(21) Application number: **91919435.7**

(22) Date of filing: **08.11.1991**

(51) Int Cl.<sup>6</sup>: **C07K 7/23, A61K 38/24**

(86) International application number:  
**PCT/EP91/02110**

(87) International publication number:  
**WO 92/08733 (29.05.1992 Gazette 1992/12)**

(54) **LHRH-ANTAGONISTS**

**LHRH-ANTAGONISTEN**

**ANTAGONISTES DE L'HORMONE DE LIBERATION DE LA LUTEOSTIMULINE (LHRH)**

(84) Designated Contracting States:  
**AT BE CH DE DK ES FR GB GR IT LI LU NL SE**

(30) Priority: **10.11.1990 CN 90108955**

(43) Date of publication of application:  
**13.10.1993 Bulletin 1993/41**

(73) Proprietor: **ASTA Medica Aktiengesellschaft  
D-01277 Dresden (DE)**

(72) Inventor: **XIAO, Shaobo,  
Tianjin Municipal Research Institute  
Nankai Dist, Tianjin 300191 (CN)**

(56) References cited:  
**EP-A- 0 277 829**

- **ENDOCRINE REVIEWS**, vol. 7, no. 1, The Endocrine Society; M.J. KARTEN et al., pp. 44-66
- **ZEITSCHRIFT FÜR NATURFORSCHUNG**, vol. 41c, 1986, Tübingen (DE); K. FOLKERS et al., pp. 1087-1091
- **ZEITSCHRIFT FÜR NATURFORSCHUNG**, vol. 42b, 1987, Tübingen (DE); K. Folkers et al., pp. 101-106
- **R.W. ROESKE et al.**, "LhRh Antagonists With Low Histamine Releasing Activity", B.H. Vickery et al., (eds.), MTP Press, Boston, MD (US), 1987; pp. 17-24
- **SCIENCE IN CHINA**, vol. 34, no. 2, February 1991, series B; K.L. LIU et al., pp. 201-208
- **INTERNATL. JOURNAL OF PEPTIDE & PROTEIN RESEARCH**, vol. 35, no. 2, February 1990, Copenhagen (DK); K.L. LIU et al., pp. 157-160

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

**EP 0 564 466 B1**

## Description

The present invention relates to novel peptides and their derivatives having exact chemical structure. The invention is also directed to the methods of their preparations and applications. Hypothalamic luteinizing hormone releasing hormone (LHRH) acts on the anterior pituitary gland to stimulate the secretion of luteinizing hormone (LH) and follicular stimulating hormone (FSH). The antagonistic analog of LHRH acts on anterior pituitary rapidly, lasts a long duration, can be safely and reversibly used for contraception or selectively suppression of gonadotropin secretion. For such kind of application, LHRH antagonists are superior to agonists. Up to now, there are more than two thousands of LHRH analogs have been designed and synthesized.

K. Liu et al., *Int. J. Peptide Protein Res.*, 35 (2), pg. 157-160 (1990) disclose LHRH antagonists, wherein the amino acid in position 6 of LHRH is exchanged against derivatives of phenylalanine.

M.J. Karten et al., *Endocrine Reviews*, 7 (1), pg. 44-66 (1986), describe GnRH analogs with modifications in position 2, 3, 6 or 10 and state that a mere increase in the number of amino acid substitutions doesn't necessarily lead to a higher potency.

K. Folkers et al., *Z. Naturforsch.*, 41C, pg. 1087-1091 (1986), and *ibid.*, 42B, pg. 101-106 (1987), teach certain modifications of LHRH.

R.W. Roeske et al., in B.H. Vickery and J.J. Nestor (Eds.), "LHRH Analogs: Contraceptive and Therapeutic Application", MTP Press, Boston 1987, pg. 17-24, disclose LHRH analogs with substitutions by MePal and/or Arg in position 5 and 6, respectively.

EP-A-0 277 829 describes LHRH analogs, wherein the arginine in Position 8 is alkylated on the side chain amino group.

The above citations show that only slight modifications in the protein sequence of LHRH lead to rather drastic changes in pharmacological properties.

"Nal-Arg" analog showed fairly high antifertility activity. However, "Nal-Arg" analog showed also very strong histamine-releasing activity (HRA). It caused transient edema of the face and extremities in rats when administered at a dosage as high as 50-100 times of therapeutic dose. The result of clinical trial demonstrated histamine-related systemic effects. Other LHRH antagonists containing DArg<sup>6</sup> or DLys<sup>6</sup> showed similar side effects, their ED<sub>50</sub> for HRA were below 1 µg/ml. The present invention provides new LHRH antagonists which have very high antioviulatory activity (AOA) and very low histamine-releasing activity (HRA) and negligible side effects. The contents and examples of this invention are as follows:

The design methodology of this invention is based on the topological similarity between the molecule of parent compound [Nac-D2Nal<sup>1</sup>, DpClPhe<sup>2</sup>, D3Pal<sup>3</sup>, Ser<sup>4</sup>, Tyr<sup>5</sup>, DArg<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]NH<sub>2</sub> (II) and a neuropeptide, Substance P, which features the modification of both alkaline and lipophilic area in the molecular of the parent compound to give new antagonists having both high AOA and low HRA. The term "modification" hereof is adjusting or substitution of the amino acids in the area of Tyr<sup>5</sup>-DArg<sup>6</sup>-Arg<sup>8</sup> in C-terminus and the aromatic acids in N-terminus of (II). More specifically, the design is introduction of suitable alkaline group and substitutions of unnatural amino acids in position 2, 3, 5, 6, 8 of (II).

The following are also the methods and examples of this invention.

1. Substitution of D3pal which is an aromatic amino acid having suitable basicity for DArg<sup>6</sup> in (II) to obtain analog (III): [Nac-D2Nal<sup>1</sup>, DpClPhe<sup>2</sup>, D3Pal<sup>3</sup>, Ser<sup>4</sup>, Tyr<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]NH<sub>2</sub>

2. Substitution of Arg<sup>5</sup> for Tyr<sup>5</sup> in (III) to obtain (IV): [Nac-D2Nal<sup>1</sup>, DpClPhe<sup>2</sup>, D3Pal<sup>3</sup>, Ser<sup>4</sup>, Arg<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]NH<sub>2</sub>

3. Substitution of Dphe<sup>3</sup> or its derivatives DXCH<sub>2</sub>Phe for D3pal<sup>3</sup> in (IV) to obtain (V): [Nac-D2Nal<sup>1</sup>, DpClPhe<sup>2</sup>, DPhe<sup>3</sup>, Ser<sup>4</sup>, Arg<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]NH<sub>2</sub> or its (DXCH<sub>2</sub>Phe<sup>3</sup>) analogs.

4. Substitutions of Dphe<sup>3</sup> or its derivatives for D3pal<sup>3</sup> in (III) to obtain (V'): [Nac-D2Nal<sup>1</sup>, DpClPhe<sup>2</sup>, DPhe<sup>3</sup>, Ser<sup>4</sup>, Tyr<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]NH<sub>2</sub> or its (DXCH<sub>2</sub>Phe<sup>3</sup>) analogs.

A series of new LHRH antagonists of the formula [Nac-D2Nal<sup>1</sup>, AA<sup>2</sup>, AA<sup>3</sup>, Ser<sup>4</sup>, AA<sup>5</sup>, AA<sup>6</sup>, Leu<sup>7</sup>, AA<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]NH<sub>2</sub> have been synthesized, where AA are natural or unnatural amino acids which are expressed as D- or L-ArAla. More specifically herein,

AA<sup>2</sup> = D-pClPhe, D-ArAla, DPhe, Ar-Ala, DXCH<sub>2</sub>Phe;

AA<sup>3</sup> = D3Pal, Ar-Ala, D-ArAla, DPhe, D-XCH<sub>2</sub>Phe;


AA<sup>5</sup> = Arg, DMap, Pip, Tyr, Pal, Mop, Tep, Map, Phe, Eap, Pap, Bap, DMop;

$$AA^6 = D3Pal, D\text{-}Ar\text{-}Ala, D\text{-}XCH^2Phe;$$


AA<sup>8</sup> = Pip, Mop, Tep, Map, Eap, Pap, Bap, Arg;

in which


Ar =



and



in which

$$X =$$


and  $R'_2N-$ ,  $RR'_1N-$

in which

$$R' = \text{CH}_3-, \text{CH}_3\text{CH}_2-, \text{C}_3\text{H}_7-, \text{C}_4\text{H}_9-, \text{H}-;$$

$R = \text{CH}_3-, \text{CH}_3\text{CH}_2-, \text{C}_3\text{H}_7-, \text{C}_4\text{H}_9-, \text{H}-;$

The LHRH antagonists obtained by using the above described method, as a kind of peptide medicine, can be used to treat the disorder of reproductive endocrine system, such as edometriosis, precocious puberty of children and to treat prostate cancer and breast cancer as well as used as male or female contraceptives for birth control, or used in the diagnosis and treatment of infertility, etc. Such peptide medicine can be prepared as normal injection injectable capsules or other formulations for real application.

Further description of this invention is as follows:

In the natural course of histamine releasing in the body, neuropeptide substance P plays a very important role, its  $\text{ED}_{50}$  for HRA is 5-15  $\mu\text{M}$ . The chemical structure of SP is  $[\text{Arg}^1, \text{Pro}^2, \text{Lys}^3, \text{Pro}^4, \text{Gln}^5, \text{Gln}^6, \text{Phe}^7, \text{Phe}^8, \text{Gly}^9, \text{Leu}^{10}, \text{Met}^{11}]\text{NH}_2$ . The study on the relationship between its structure and HRA showed that  $\text{Arg}^1\text{-Pro}^2\text{-Lys}^3$  in the N-terminus in the molecule of SP is essential for its HRA because deletion of these three amino acids from the molecule entirely abolished its HRA. By contrast, deletion of one two or three amino acids in C-terminus remained HRA as high as one fourth as HRA of itself. Further deletion of  $\text{Phe}^8$  and  $\text{Phe}^7$ , HRA reduced 4 % and 0,57 % of those of SP. Further deletion of  $\text{Gln}^{5,6}$  did not cause significant change of HRA. The above data implies that the lipophilic area around  $\text{Phe}^{7,8}$  determines the value of HRA, this area involves in the binding of molecule with the receptor of mast cell.

As mentioned previously,  $(\text{D2Nal}^1, \text{DArg}^6)$  analogs of LHRH showed very high HRA, its molecular structure has topological similarity with SP:  $\text{DArg}^6\text{-Leu}^7\text{-Arg}^8$  in the former appears to be corresponding to  $\text{Arg}^1\text{-Pro}^2\text{-Lys}^3$  in the latter, both consist of a pair of strongly basic amino acid residues between which only one neutral amino acid residue is present, i. e. both  $[\text{D2Nal}^1, \text{DArg}^6]$  analog of LHRH and SP contains two strongly basic amino acid residues which are in 1,3 position relationship. On the other hand, a cluster of aromatic amino acid residues in the former is considered to be corresponding to  $\text{Phe}^{7,8}$  area in SP in terms of determination of the magnitude of HRA.

The design of this invention consists of two aspects: one is modifying  $\text{Tyr}^5\text{-DArg}^6\text{-Arg}^8$  area in C-terminus, the other is fine adjusting the aromatic acids after optimizing the modification of the alkaline area in C-terminus.  $[\text{Nac-D2Nal}^1, \text{DpClPhe}^2, \text{D3Pal}^3, \text{Ser}^4, \text{Tyr}^5, \text{DArg}^6, \text{Leu}^7, \text{Arg}^8, \text{Pro}^9, \text{DAla}^{10}]\text{NH}_2$  (II) is used as parent compound, which showed AOA 100 % at 0,5  $\mu\text{g}$  in corn oil, 57 % at 0,25  $\mu\text{g}$ .

First,  $\text{DArg}^6$  in (II) could be replaced by weakly basic or neutral aromatic acids, such as  $\text{D3Pal}$ ,  $\text{D6Qal}$ , tetrahydrotryptophan, methyl tryptophan.  $[\text{Nac-D2Nal}^1, \text{DpClPhe}^2, \text{D3Pal}^{3,6}, \text{DAla}^{10}]\text{LHRH}$  (III) was obtained when  $\text{D3Pal}^6$  was substituted for  $\text{DArg}^6$  in (II). (III) showed AOA 100 % at 3  $\mu\text{g}$ , 83 % at 1  $\mu\text{g}$  (in corn oil), and its  $\text{ED}_{50}$  for HRA was 9.8  $\mu\text{g}/\text{ml}$ , much better than that of "Nal-Arg" analog  $\text{ED}_{50}$  for HRA was less than 1  $\mu\text{g}/\text{ml}$ . It seems that the basicity of the whole molecule should equal to or closed to that of a pair of arginine in order to obtain high AOA. Because position 5, like position 6, does not involve in the receptor binding, a wide variety of amino acid including arginine can be inserted in position 5. A series of new analogs were designed. For example, substitution of  $\text{Arg}^5$  for  $\text{Tyr}^5$  in (III) gave  $[\text{Nac-DNal}^1, \text{DpClPhe}^2, \text{D3Pal}^{3,6}, \text{Arg}^5, \text{DAla}^{10}]\text{LHRH}$  (IV). Both (IV) and (II) contained two arginines, but the distance between two arginines in (IV), whose geometric relationship became 1, 4, i. e. there were two other amino acids between these two arginine, was larger than that in (II). Therefore, HRA would be reduced and, on the other hand, because of the presence of two arginine, AOA should not be lower than that of (II). The bioassay result of (IV) showed that  $\text{ED}_{50}$  for HRA was 3,5  $\mu\text{g}/\text{ml}$ , while AOA was 60 % at 0,12  $\mu\text{g}$  (corn oil), 85 % at 0,25  $\mu\text{g}$ , 100 % at 0,5  $\mu\text{g}$ . This was the first time for LHRH antagonists to achieve  $\text{ED}_{50}$  for AOA which was equal or less than 0,125  $\mu\text{g}$ .

Therefore further design was based on the structure of (IV).

There are four alkaline residues,  $\text{D3Pal}^{3,6}$  and  $\text{Arg}^{5,8}$  in the molecule (IV), while (II) contains only three alkaline. Therefore, it is reasonable to replace one  $\text{D3Pal}$  by a neutral amino acid; on the other hand, (IV) showed very strong hydrophilicity and reducing the hydrophilicity by the substitution of a hydrophobic amino acid for  $\text{DPal}$  in (IV) would be beneficial to the retention of the drug in the body and then to the extension of the effective duration. A new series of analogs were then designed by substitution for  $\text{D3Pal}^3$ . (V) showed 100 % of AOA at 1  $\mu\text{g}$  (in saline), equal to that of parent compound (IV), while HRA reduced by a half:  $\text{ED}_{50}$  for HRA was 7,4  $\mu\text{g}/\text{ml}$ .

Further substitution of  $\text{DPhe}^2$  for  $\text{DCIPhe}^2$  reduced the lipophilicity of this area in the molecule and reduced HRA.

$\text{Arg}^5\text{-D3Pal}^6\text{-Leu}^7\text{-Arg}^8$  in the C-terminus of (IV) seems to play a major role in triggering histamine releasing.  $\text{D3Pal}$  combines aromaticity, basicity and hydrophilicity in one molecule, it is also stereo-acceptable in LHRH antagonists for receptor binding. Similarly, design of new series of unnatural amino acids possessing the same character as  $\text{D3Pal}$  may lead to better LHRH antagonists than (IV) or (V).

Modification of natural, lipophilic, aromatic amino acid e. g. phenylalanine, for example, by means of the method described below in The Synthesis of Novel Unnatural Amino Acids, lead to a series of novel amino acids which combine aromaticity, hydrophilicity and basicity in one molecule and can be expressed as the formula:  $\text{R}_1\text{R}_2\text{NCH}_2\text{C}_6\text{H}_4\text{CH}_2\text{CH}(\text{NH})\text{CO}_2\text{H}$  (VI), where  $\text{R}_1$  and  $\text{R}_2$  may be the same or may differ each other, may be chain-like or cyclic, may also contain hetero-atom. With the  $\text{R}_1$  and  $\text{R}_2$  change, a series of amino acid can be obtained, which show systemically changed basicity, hydrophilicity and stereo-character. Introduction of those amino acids in position 5, 6, 8 of (IV) have given three series of new antagonists of LHRH. The bioassay results showed that each series gave at least one new antagonist

showing 100 % AOA at 1  $\mu$ g, similar to that of (IV), while HRA was significantly reduced. An example was (VII): [NAC-D2Nal<sup>1</sup>, DpClphe<sup>2</sup>, D3Pal<sup>3</sup>, Ser<sup>4</sup>, Mop<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]-NH<sub>2</sub>, which showed 100 % of ADA at 1  $\mu$ g, ED<sub>50</sub> 14,7  $\mu$ g/ml for HRA, appeared better than (V). When substitution of (VI) for Arg<sup>8</sup> in (IV), the extent of HRA decrease was positively correlated to the length of R in (VI), so ED<sub>50</sub> for HRA could be higher than 200  $\mu$ g/ml, such kind of compounds can be easily dissolved in aqueous solution and expected to be utilized clinically without formulation problems. The results demonstrated that Arg<sup>8</sup> or Lys<sup>8</sup> was not essential for highly potent LHRH antagonists. Suitable alkaline center in position 8 can ensure high AOA, meanwhile activity inducing mast cell to release histamine was remarkably reduced when the basic enter mentioned above possessing significant stereo-hinder.

This invention combining the modification in both N-and C-terminus lead to better LHRH antagonists.

The process of synthesis are illustrated as follows:

#### 1. The synthesis of Novel Unnatural Amino Acids


Over 60 series and nonseries, D- or L-amino acids are designed and synthesized through the four synthetic routes outlined in the schema below. The structure of these unnatural amino acids are shown with the general structural formulas listed in the same schema. Some of these amino acids have alkalinity, hydrophilicity and aromaticity respectively, while the others have them all together in the molecule.




Path 1

Path 2


5



10




15




20


25




30



35




40







45

50



55



5

10

15

20

25

30


35




40

45


50

55






5




10




15


20



25



30




35

40



45

50



55





Path III





Chymotrypsin
















Secondary  
Amine







SOCl<sub>2</sub>/EtOH







Path IV D or L  $\text{Ar}-\text{CH}_2-\underset{\text{NH}_2}{\text{CH}}-\text{COOH}$ , wherein



## 2. Synthesis of Peptide

The synthesis begins from the C-terminus of the peptide on benzhydrylamine hydrochloride resin (BHA resin) utilizing the method of solid-phase peptide synthesis introduced by Merrifield. It is a three-step process including anchor, coupling and cleavage. Dichloromethane (DCM) is the major solvent used for washing between each step of reaction while isopropanol alcohol (IPA) and N,N-dimethylformamide (DMF) are also used when it is necessary. Catalyzed by excessive dicyclohexylcarbodiimide (DCC), coupling reaction is carried out, while adequate amount of 1-hydroxybenzotriazole (HOBT) is added. The degree of the coupling reaction is monitored with Kaises ninhydrin method. The second coupling reaction would be carried out if it gives a positive result in Kaises test. The peptide chain is cleaved from the resin using anhydrous hydrogen fluoride (HF) in the presence of anisole after the completion of all reactions necessary



on the resin, all of the temporary protecting group are deprotected at the same time. After washed by ethyl acetate or ether crude products of LHRH antagonists are obtained by aqueous acetic acid extraction followed by lyophilization. The yield is over 50 %.

### 5 3. Purification of Peptide

(1) The peptide is purified by gel permeation chromatography or silica partition chromatography through a column as high as 60 - 100 cm with the aid of UV/TLC monitoring. The LHRH antastits purified once are obtained after lyophilizing the major fractions. The yield is 50 - 90 % and the purity can be over 90 %.

10 (2) The peptide then is further purified on Waters high performance liquid chromatography (HPLC) instrument using reverse phase C18 column (7,8 x 300 mm) ( $\mu$ -Bondapak 84176). The yield of this step is 20-50 % while the purity is no less than 99 %.

### 15 4. Purity Analysis of Peptide

#### (1) TLC analysis

It is carried out on a plastic sheet coated by silica gel 60 F254 of 5 - 10 cm height. They all shows a single spot when developed in five different solvent systems.

#### 20 (2) HPLC analysis

They all shows a single peak when eluted with two kinds of solvent system, respectively, utilizing Waters HPLC instrument on a analytic column ( $\mu$ -Bondapak 27324) when monitored by UV 210. The sample size is 10 - 200  $\mu$ g.

### 25 5. Amino Acid Analysis of Peptide:

According to the PICO-TAG method developed by Waters Company, 50  $\mu$ g of sample which have been dried under vacuum over 2 hours is weighed accurately on a  $10^{-5}$ g scale balance. After dissolved in water, 10  $\mu$ g of aliquot is added to a reaction tube in which 1:1 hydrochloride acid (containing 1 % phenol) was added according to the manual.

30 The reaction lasts 22-24 hours at 105°C in a sealed container which had been filled with nitrogen and pumped to vacuum to remove the oxygen in reaction tube. Phenol isothiocyanate is added to derive the amino group after evaporating of excessive hydrochloride acid. Then it was analyzed with the HPLC-instrument equipped with PICO-TAG amino acid analytical column and monitored by UV254. The content of each amino acid and the relative mole ratio were calculated to give the amino acid composition of the sample based on the comparison of the integrated area of each amino acid to that of H-standard sample of Waters. The classical ion-exchange-ninhydrin derivation method (IEN) was also used as control which gave the same results. But it needed ten times more sample to get a satisfied result.

### 40 6. Evaluation of biological activity:

Corbin's rat antioovulation method is used. The healthy, adult, female SD rats (BW 200-250 g) are used in this experiment. All animals are maintained at 22-24 °C and on 14 h/10 h (light/dark) schedule. They are given standard food, and water ad libitum. The rats showing at least two consecutive 4-day estrous cycles in vaginal smear examination can be used in this experiment. The rats are given peptides (LHRH antagnists) at noon of proestrous with different dose in saline solution. The rats are sacrificed next day, their oviduct of two sides are exammed under a dissecting microscope to determine the ovum number. The rats were divided into several groups according to the dosing, each group consists of about 10 rats, and the control group in which the rats are given equal amount of saline consists of 9-10 rats. The antioovulatory activity (AOA) is shown in the following equation:

$$50 \quad \text{AOA} = \frac{\text{number of unovulated rats}}{\text{Total number of treated rats}} \times 100 \%$$

### 7. Evaluation of Histamine Releasing Activity:

#### 55 (1). Histamine releasing test (HRT) in vitro:

The healthy, adult, male SD rats (BW 200-250 g) housing in the above same conditions are used in this experiment. After anesthetized by CO<sub>2</sub> the peritoneal cavity is washed with 50 ml of PIPES AC medium containing 20 units of

heparin. Following centrifugation at 200xg for 8 min at 4 °C, cells are washed again and finally resuspended to a concentration of  $8 \text{ to } 24 \times 10^5$  total leucocytes/ml in PIPAS AC. This suspension contains approximately 5-10 % mast cells. Washed cells are used immediately after collection and are prewarmed for 5 min at 37 °C prior to pipetting 0,3 ml aliquots into polystyrene tubes containing 0,3 ml of diluted peptide. The mixtures are incubated for 15 min at 37 °C and the reaction stopped by centrifugation at 400 xg for 15 min at 4 °C. The supernatans are assayed for histamine content by manual fluorometric assay method after successively extraction with n-butanol and n-heptane. The histamine content can be obtained from the histamine standard curve (see below). The percentage of histamine release can be calculated from the following equation:

$$\text{Histamine release (\%)} = \frac{E-B}{C-B} \times 100 \%$$

where E is the fluorometric reading of experimental sample, B is the fluorometric reading of samples with cells and buffer only, and C is the fluorometric reading of "complete" (cells treated with  $\text{HClO}_4$ ).

The standard curve can be obtained by plotting the OD values on a fluorometer at 350 nm/450 nm (activation/fluorescent) against the concentrations of serially diluted solution of accurately weighted histamine hydrochloride. The relative parameter r of the histamine standard curve can be 0.9998, and the lowest detectable concentration of histamine is 0,5 ng/ml.

The  $\text{ED}_{50}$  value of peptide can be gotten from the dose response curves obtained by plotting the histamine release versus the peptide concentration on semilogarithmic paper.

All peptide samples should be tested with mast cells from a minimum of 3 different rats.

## (2). Cutaneous anaphylactoid activity test (CAT):

The healthy, adult, female SD rats (BW 250 g) are used in this experiment. The rats are injected intravenously with Evan's blue (1ml of 0,05 % solution). Immediately after that, the 0,05 ml of peptide solution (5, 0,5 and 0,05  $\mu\text{g}/\text{ml}$  respectively) and saline (control) are injected intradermally into a shaved section on the back of the animals. 30 minutes after the injection, the rats are sacrificed and the dorsal skin was reflected. The diameters of the lesions are measured in millimeters in two perpendicular directions with a vernier caliper. The diameter of control is usually less than 5,5 mm.

The amount of Evan's blue permeating into the skin from the blood vessel can be spectrophotometrically measured, too. The skin corresponding to the lesion area is cut down and immersed in a mixture of acetone/saline (7:3; Vol/Vol) overnight. After centrifugation next day, the content of Evan's blue in the supernatant is measured with a spectrophotometer (UV-260) at 610 nm against reference solution of acetone/saline (7:3). Each peptide were tested in a minimum of 3 different rats.

A variety of new LHRH antagonists were designed and synthesized by means of the method described above. In brief, the new structure of LHRH antagonists was obtained by single or multiple substitution of the various natural and unnatural amino acids listed in the previous paragraphs.

A part of examples of new LHRH antagonists obtained thereupon are illustrated in table 1

55

Analogue	AA 1	AA 2	AA 3	AA 4	AA 5	AA 6	AA 7	AA 8	AA 9	AA 10
Parent	NAC-D2NADpClPhe	D3Pal	Ser	Tyr	DArg	Leu	Arg	Pro	DAla-NH2	
				Arg						
				Arg	DPhe			P1p		
				Arg	DMop			P1p		
				Arg	DPhe			Mop		
				Mop	D3Pal					
		DP1p		Mop	DMop			P1p		
				Arg	D3Pal			Pap		
				Arg	D3Pal			P1p		
				DFPhe	D3Pal			Pap		
					DP1p			Eap		
					DMap			Mop		
		DPhe		Arg	DMop			Map		
		OPClPhe			DP1p			Map		
		DPhe		Arg	D3Pal					
				Eap				Mop		
				Tep	DMop			Pep		
				Tep	DMap			Mop		
				Tep	DEap					
				Tep	DBap					
				Tep	DPap					
				Tep	DTep					
					DMop			Mop		
	DpFPhe	D3Pal								
	DPhe			Mop	DMop			Eap		
					DP1p			Pap		

Table 1: A Part of Examples Related to This Invention, Page 2

Analogue	AA 1	AA 2	AA 3	AA 4	AA 5	AA 6	AA 7	AA 8	AA 9	AA 10
Parent	NAC-D2NaDpClPhe		D3Pal	Ser	Tyr	DArg	Leu	Arg	Pro	DAla-NH2
						DBap		Pip		
					Tep	DMop		Eap		
						DMop				
						DTep				
					Tep	DMop				
						DTep				
		DPClPhe								
		DPClPhe				DMop		Pip		
						DMop		Bap		
					Mop					
					Mop	D3Pal				
					Arg	DMop				
					Arg	DPip				
					Arg	DTep				
						D3Pal		Pip		
					Arg	D3Pal		Pip		
					Arg	D3Pal		Mop		
					Arg	D3Pal		Tep		
					Arg	D3Pal		Pap		
					Arg	DTep		Pip		
					Arg	DTep		Mop		
					Arg	DTep		Tep		
		DPhe			Arg	DTep		Mop		
		DFPhe			Arg	DTep		Mop		
		DFPhe			Arg	DTep		Eap		
		DFPhe			Arg	DTep		Pap		
		DFPhe			Arg	DTep		Bap		
		Nep			Tep	DPap				

### The Applications of This Invention:

1. After finishing the preclinical pharmacology and toxicology study, we can apply these new LHRH antagonists which have high therapeutic effectiveness and low side effect in clinic so as to develop new medicine for treating the endometriosis and their disorder in reproductive endocrine system including precious puberty of children, prostate cancer and breast cancer. Since they suppress the secretion of gonadotropin through competing receptor with endogenous LHRH, and act rapidly reversibly and safely, they can be further developed as new type of contraceptives for male or female. Besides, they can be also used in treatment of infertility and for selectively and reversibly abolishing the function of pituitary gland in terms of secreting gonadotropin.

Being a kind of peptide medicine, the LHRH antagonists described herein are unlikely to be administered orally.

But they can be easily made into lyophilized powder which are ready to dissolve in saline for injecting iv, sc or im.

Moreover, long-actin delivery systems, such as biodegradable, injectable capsules are studied. The capsules can be implanted subcutaneously by a special syringe and would be adsorbed by the tissue after release of all peptide contents and do not need to remove surgically. The long-acting delivery system is specially useful for long-term administration of LHRH analogues in clinic.

The following are the analyses results of the examples (taking three analogues IV, V, VII as typical examples):

#### (1) The Purity

Thin layer chromatography (TLC):

There is only a single spot in each of the chromatogram developed in five different solvent systems.

High performance liquid chromatography (HPLC):

There is only a single peak in each of the chromatogram eluted with two different solvent systems.

The values of Rf and retention time TR are shown in Table 2, also with reference of Figure 1-4.

Table 2:

The chromatographic analysis results of LHRH antagonists							
Analogues	HPLC		TLC				
	TR1	TR2	Rf1	Rf2	Rf3	Rf4	Rf5
		(min)					
IV	7.55	5.26	0.23	0.21	0.31	0.19	0.65
V	7.90	8.11	0.32	0.30	0.35	0.30	0.69
VII	16.19	9.58	0.17	0.08	0.16	0.40	0.12

Solution A + 80 % acetonitrile

Solvent System 2:

Solution A is 0.01M  $\text{KH}_2\text{PO}_4$  aqueous solution (pH3)

Solution B is 20 % solution A + 80 % acetonitrile

TLC solution system:

1. nBuOH/EtOAc/HOAc/ $\text{H}_2\text{O}$  (5:5:1:1)
2. nBuOAc/nBuOH/HOAc/ $\text{H}_2\text{O}$  (2:8:2:3)
3. nBuOAc/HOAc/ $\text{H}_2\text{O}$  (4:1:5), up phase
4. nBuOH/HOAc/ $\text{H}_2\text{O}$  (4:1:2)
5. nBuOH/EtOAc/HOAc/ $\text{H}_2\text{O}$  (1:1:1:1)

#### (2) Amino acid analysis

The analysis are carried out according to the method of classical IEN and new Pico-Tag, the results are shown in Table 3 and Figure 5,6.

Table 3:

The amino acid composition of LHRH antagonists										
Analogues	Methods	Ser	Arg	Ala	Pro	Leu	Phe	Pal	pCIPhe	Nal
IV	IEN	0.86	2.05	1.01	0.99	1.13		+	+	ND
	Pico-TAG	0.92	2.25	0.91	1.01	0.91		+	+	+
V	IEN	0.81	2.02	1.03	1.03	0.12	0.99	+	+	+
	Pico-TAG	0.68	2.26	0.93	1.29	1.04	1.00	+	+	+
VI	IEN	0.91	0.91	1.00	1.00	1.09		+	+	ND
ND: Not determined										

## (3) The bioassay results

The results of bioassays including antioviulatory activity at different doses and ED<sub>50</sub> for histamine-realising activity in vitro are illustrated in table 4 in which 26 antagonists are listed as examples.

Table 4:

Bioassay Results of New LHRH Antagonists based on Parent structure							
	Substituted Amino Acids	% AOA/ $\mu$ g					HRA ( $\mu$ g/ml)
		0.125	0.25	0.5	1.0	2.0	ED <sub>50</sub> $\pm$ SEM
1	Parent		50	75	100		3.5 $\pm$ 0.38
2	DPhe		29	60	100		7.4 $\pm$ 0.98
3	DPhe, DPhe				0		18.5 $\pm$ 7.00
4	DTyr, Lys				40		5.1 $\pm$ 2.15
5	D-Phe					60	35.0 $\pm$ 5.05
6	Map			29			24.8 $\pm$ 4.47
7	Eap			43			12.0 $\pm$ 0.50
8	Pap			0			9.6 $\pm$ 0.19
9	Bap			14			23.5 $\pm$ 5.78
10	D-Map			12,5			18,3 $\pm$ 2,38
11	Tep			14			36.8 $\pm$ 5.68
12	Pip	17	33	71	100		9.4 $\pm$ 1.63
13	Mop			25	100		14.7 $\pm$ 2.70
14	D-Map			14			19.5 $\pm$ 2.50
15	D-Eap			14			13.0 $\pm$ 1.00
16	D-Tep			71			22.5 $\pm$ 3.25
17	D-Pip			0	50	57	7.6 $\pm$ 2.48
18	D-Mop		33	67	100		> 11
19	Map			57	100		5.4 $\pm$ 1.22
20	Eap				29		56.9 $\pm$ 15.1
21	Pap				50	88	70.4 $\pm$ 26.8
22	Bap				0		> 235
23	Tep				100		6.6 $\pm$ 2.13
24	Pip				43		27.5 $\pm$ 2.50
25	Mop				71		52.5 $\pm$ 17.5
26	D-Map				0		28.0 $\pm$ 9.00

\* The parent structure is: [NAc-D<sup>2</sup>Nal<sup>1</sup>, DpClphe<sup>2</sup>, D<sup>3</sup>Pal<sup>3</sup>, Ser<sup>4</sup>, Arg<sup>5</sup>, D<sup>3</sup>Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]NH<sub>2</sub>

As illustrated and described above, the LHRH antagonists designed and synthesized according to this invention shows very good properties. They are pure in TLC or HPLC analysis. Their compositions are correct, i. e., the same is designed. Their antifertile activity is high: they can inhibit rat ovulation when injected s. c. at the dosage of 0.1 to 2.0  $\mu$ g at the noon of proestrus. Their histamine related side effect is low: their ED<sub>50</sub> for in vitro histamine releasing activity (the effective dose for rat mast cell to release 50 % of histamine) is ranged 5-300  $\mu$ g/ml; the lesion induced by them in the cutaneous anaphylactoid test in rats is as small as required in clinic. Their water-solubility is very good, all bioassays are carried out in saline solution, so they are easy to formulate for injection in clinic. They are also ready to be formulated as long-acting delivery systems, among which injectable microcapsules are most convenient for long-term suppression of gonadotropin and gonadal hormone. Therefore, they can be used as highly effective, reversible and safe contraceptives for both male and female. They can be also utilized for treatment of various diseases related to disorders of reproductive endocrine such as hormone-dependent prostate cancer and breast cancer, endometriosis, precocious puberty of children. They are also useful in treatment of infertility. The new LHRH antagonists herein can also be utilized in the basic research of reproductive physiology and pharmacology, such as in the study on the function of pituitary gland, on the effect of gonadal hormones or gonadotropins or LHRH on sexual behaviour, etc.

ABBREVIATIONS

The following are abbreviations which have been used in the text of this patent application document.

5	Ala	alanine	
	AOA		antioviulatory activity
	Arg		arginine
10	Bap		dibutylaminomethyl phenylalanine
	Boc		t-butyloxy carbonyl
	BuOAC	butyl acetate	
15	CAT		cutaneous anaphalactoid test
	DCC		dicyclohexylcarbodiimide
	DCM		dichloromethane
20	D2Na1	D- $\beta$ -(2-naphthyl) alanine	
	D3Pa1	D- $\beta$ -(3-pyridyl) alanine	
	DpClPhe	p-chloro-D-phenylalanine	
	DpFPhe	p-fluoro-D-phenylalanine	
25	D6Qa1	D- $\beta$ -(6-quinolyl) alanine	
	DMF		N,N-dimethyl formamide
	Eap		diethylaminomethyl phenylalanine
30	ED <sub>50</sub>	effective dose for 50 % response	
	EtOAC	ethyl acetate	
	FSH		follicle-stimulating hormone
35	Glu		glutamic acid
	Gly		glycine
	His		histidine
	HOBT	1-hydroxybenzotriazole	
40	HPLC	high performance liquid chromatography	
			ninhydrin derivation
	HRA		histamine-releasing activity
45	HRT		histamine-releasing test
	IEN		ion exchange chromatography with
			post-column
50			
55			

	IPA	isopropyl alcohol
	LH	lutinizng hormone
5	LHRH	lutinizng hormone releasing hormone
	Leu	leucine
	Lys	lysine
	Map	dimethylaminomethyl phenylalanine
10	Met	methionine
	Mop	mophorlinomethyl phenylalanine
	nBuOH	n-butyl alcohol
15	NS	normal saline
	Pap	dipropylaminomethyl phenylalanine
	Phe	phenylalanine
20	Pip	piperidinomethyl phenylalanine
	Pipes	piperazine-N,N'-bis[2-ethanesulfonic acid]
	Pro	proline
	Rf	rate of flow
25	SE	standard error
	Ser	serine
	TFA	trifluoroacetic acid
30	TLC	thin-layer chromatography
	TR	retention time
	Trp	tryptophan
35	Tyr	tyrosine
	Tep	tetrahydropiperidyl methyl phenylalanine

40

**Claims**

1. [Nac-D2Nal<sup>1</sup>, DpCIPhe<sup>2</sup>, DPhe<sup>3</sup>, Ser<sup>4</sup>, Arg<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]NH<sub>2</sub>
- 45 2. [Nac-D2Nal<sup>1</sup>, DpCIPhe<sup>2</sup>, DPhe<sup>3</sup>, Ser<sup>4</sup>, Tyr<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]NH<sub>2</sub>
3. [N-AC-D2Nal<sup>1</sup>, P-CI-D-Phe<sup>2</sup>, D3Pal<sup>3</sup>, Ser<sup>4</sup>, Mop<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, D-Ala<sup>10</sup>]NH<sub>2</sub>
4. [N-AC-D-2Nal<sup>1</sup>, D-Phe<sup>2</sup>, D3Pal<sup>3</sup>, Ser<sup>4</sup>, Mop<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, D-Ala<sup>10</sup>]NH<sub>2</sub>
- 50 5. [N-AC-D-2Nal<sup>1</sup>, P-CI-D-Phe<sup>2</sup>, D3Pal<sup>3</sup>, Ser<sup>4</sup>, Arg<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Pap<sup>8</sup>, Pro<sup>9</sup>, D-Ala<sup>10</sup>]NH<sub>2</sub>
6. The use of one of the LHRH antagonists of claims 1 to 5 for the production of a medicament for the treatment of disorders of the reproductive endocrine system and prostate and breast cancer.
- 55 7. The use of one of the LHRH antagonists of claims 1 to 5 for the production of a medicament for the use as contraceptive as well for diagnosis and treatment of infertility.



8. A method for producing the LHRH antagonists of claims 1 to 5 wherein the alkalinous and lipophilic areas of [NAC-D2Nal<sup>1</sup>, DpCIPhe<sup>2</sup>, D3Pal<sup>3</sup>, Ser<sup>4</sup>, Tyr<sup>5</sup>, Arg<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]NH<sub>2</sub> are modified by the replacement of the respective amino acids.

5

# Patentansprüche

1. [NAC-D2Nal<sup>1</sup>, DpCIPhe<sup>2</sup>, Dphe<sup>3</sup>, Ser<sup>4</sup>, Arg<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]NH<sub>2</sub>
- 10 2. [NAC-D2Nal<sup>1</sup>, DpCIPhe<sup>2</sup>, DPhe<sup>3</sup>, Ser<sup>4</sup>, Tyr<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]NH<sub>2</sub>
3. [NAC-D2Nal<sup>1</sup>, P-CI-D-Phe<sup>2</sup>, D3Pal<sup>3</sup>, Ser<sup>4</sup>, Mop<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, D-Ala<sup>10</sup>]NH<sub>2</sub>
4. [NAC-D-2Nal<sup>1</sup>, D-Phe<sup>2</sup>, D3Pal<sup>3</sup>, Ser<sup>4</sup>, Mop<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, D-Ala<sup>10</sup>]NH<sub>2</sub>
- 15 5. [NAC-D-2Nal<sup>1</sup>, P-CI-D-Phe<sup>2</sup>, D3Pal<sup>3</sup>, Ser<sup>4</sup>, Arg<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Pap<sup>8</sup>, Pro<sup>9</sup>, D-Ala<sup>10</sup>]NH<sub>2</sub>
6. Der Gebrauch von einem der LHRH-Antagonisten nach den Ansprüchen 1 bis 5 zur Herstellung eines Medikaments für die Behandlung von Störungen des endokrinen Fortpflanzungssystems sowie von Prostata- und Brustkrebs.
- 20 7. Der Gebrauch von einem der LHRH-Antagonisten nach den Ansprüchen 1 bis 5 zur Herstellung eines Medikaments zur Verwendung als Empfängnisverhütungsmittel sowie als Mittel zum Diagnostizieren und Behandeln der Unfruchtbarkeit.
- 25 8. Ein Verfahren zur Herstellung der LHRH-Antagonisten nach den Ansprüchen 1 bis 5, bei dem die alkalischen und lipophilen Bereiche von [Nac-D2Nal<sup>1</sup>, DpCIPhe<sup>2</sup>, D3Pal<sup>3</sup>, Ser<sup>4</sup>, Tyr<sup>5</sup>, Arg<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]NH<sub>2</sub> durch Substituierung der entsprechenden Aminosäuren modifiziert sind.

30


# Revendications

1. [Nac-D2Nal<sup>1</sup>, DpCIPhe<sup>2</sup>, Dphe<sup>3</sup>, Ser<sup>4</sup>, Arg<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]NH<sub>2</sub>.
- 35 2. [Nac-D2Nal<sup>1</sup>, DpCIPhe<sup>2</sup>, DPhe<sup>3</sup>, Ser<sup>4</sup>, Tyr<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]NH<sub>2</sub>
3. [N-AC-D2Nal<sup>1</sup>, P-CI-D-Phe<sup>2</sup>, D3Pal<sup>3</sup>, Ser<sup>4</sup>, Mop<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, D-Ala<sup>10</sup>]NH<sub>2</sub>
4. [N-AC-D-2Nal<sup>1</sup>, D-Phe<sup>2</sup>, D3Pal<sup>3</sup>, Ser<sup>4</sup>, Mop<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, D-Ala<sup>10</sup>]NH<sub>2</sub>.
- 40 5. [N-AC-D-2Nal<sup>1</sup>, P-CI-D-Phe<sup>2</sup>, D3Pal<sup>3</sup>, Ser<sup>4</sup>, Arg<sup>5</sup>, D3Pal<sup>6</sup>, Leu<sup>7</sup>, Pap<sup>8</sup>, Pro<sup>9</sup>, D-Ala<sup>10</sup>]NH<sub>2</sub>
6. Utilisation d'un des antagonistes de LHRH selon les revendications 1 à 5 pour la production d'un médicament pour le traitement des troubles du système endocrine de la reproduction et pour le traitement du cancer de la prostate et du sein.
- 45 7. Utilisation d'un des antagonistes de LHRH selon les revendications 1 à 5 pour la production d'un médicament destiné à l'emploi comme contraceptif ainsi que pour le diagnostic et le traitement de la non fertilité.
- 50 8. Un procédé pour produire les antagonistes de LHRH selon les revendications 1 à 5, caractérisé en ce que les zones alcalines et lipophiles de [Nac-D2Nal<sup>1</sup>, DpCIPhe<sup>2</sup>, D3Pal<sup>3</sup>, Ser<sup>4</sup>, Tyr<sup>5</sup>, Arg<sup>6</sup>, Leu<sup>7</sup>, Arg<sup>8</sup>, Pro<sup>9</sup>, DAla<sup>10</sup>]NH<sub>2</sub> sont modifiées par le remplacement des aminoacides respectifs.


55

Figure 1: The TLC result of LHRH antagonists IV, V, VII  
in five different systems

Sample IV



Sample V



Sample VII




Figure 2: The reversed phase HPLC spectra for the pure sample of LHRH antagonist IV

Conditions:

Column:  $\mu$ -Bondapak C18 (3.9 mm X cm)


mobile phase: A, 0,1 M  $\text{NH}_4\text{OAC}$  (pH7)

B, 20 % A + 80 % acetonitrile

gradient procedure: B from 10 % to 100 % in 15 minutes

flow rate: 2 ml/minute

detector: UV 229 nm



Column:  $\mu$ -Bondapak C18 (3,9 mm X 30 cm)

moving phase: A, 0.01 M  $\text{KH}_2\text{PO}_4$  pH 3)

B, 20 % A + 80 % acetonitrile

gradient procedure: B from 40 % to 100 % in 15 minutes

flow rate: 2 ml/minute

detector: UV 210 nm




Figure 4: The HPLC spectra for the pure sample of LHRH antagonist VII

Conditions:

Column:  $\mu$ -Bondapak C18 (3,9 mm X 30 cm)

moving phase: A, 0,01 M  $\text{KH}_2\text{PO}_4$  (pH 3)

B, 20 % A + 80 % acetonitrile

gradient procedure: B from 40 % to 100 % in 15 minutes

flow rate: 1 ml/minute

detector: UV 210 nm




Figure 5: The PICO-TAGTM spectra of LHRH antagonist IV




Figure 6: The PICO-TAGTM spectra of LHRH antagonist V

